arXiv:1301.2870v1 [math.AG] 14 Jan 2013

Kato-Usui partial compactifications over the toroidal compactifications of Siegel spaces

Tatsuki Hayama

Abstract. We construct a fan which gives a Kato-Usui partial compactification of a period domain of weight -1. This fan is given as with the fans of toroidal compactifications.

1. Introduction

Let $\mathscr{H}$ be a Siegel upper half space, and let Γ be an arithmetic subgroup. By Mumford et al. [**AMRT**] we have a toroidal compactification $\Gamma\backslash\mathscr{H}_{\Sigma_{\mathrm{tor}}}$ by taking a suitable fan Σ_{tor}. Moreover Carlson, Cattani and Kaplan [**CCK**] showed a relationship between the boundary of $\Gamma\backslash\mathscr{H}_{\Sigma_{\mathrm{tor}}}$ and the degenerating Hodge structures. These boundary points are corresponding to the nilpotent orbits, which generate limiting mixed Hodge structures by Schmid [**S**].

Generalizing this Siegel case, i.e. the case for Hodge structures of weight 1, Kato and Usui [**KU**] have introduced a construction to make partial compactifications of period domains. Let D be a period domain of pure Hodge structure introduces by Griffiths [**G**]. Then the real Lie group $G_{\mathbb{R}}$ acts on D transitively. A boundary point of Kato-Usui partial comapactifications is a nilpotent orbit generated by a nilpotent cone in $\mathfrak{g}$. Let Γ be a subgroup of $G_{\mathbb{Z}}$, and Σ be a fan of nilpotent cones which is (strongly) compatible with Γ. We can define the set $\Gamma\backslash D_{\Sigma}$ of σ-nilpotent orbits for $\sigma \in \Sigma$ modulo Γ. Here $\Gamma\backslash D_{\Sigma}$ has a log geometrical structure and is a moduli space of log Hodge structures if Γ is neat.

In Siegel case, Σ_{tor} is compatible with an arithmetic subgroup Γ and covers all possible nilpotent cones. However, beyond the Siegel case, we do not know how to construct such a big fan except for some examples. In this paper, we introduce how to construct a fan which covers all possible nilpotent cones of special type.

. We review [**H**] to clarify our result in this paper. Let D be a period domain of weight -1, i.e. odd-weight by twisting. Then $G_{\mathbb{R}}$ is isomorphic to the symplectic group. Here an isotropy subgroup L of $G_{\mathbb{R}}$ is compact. Let K be the maximal subgroup of $G_{\mathbb{R}}$ containing L. We then have the natural projection

$$D \cong G_{\mathbb{R}}/L \to G_{\mathbb{R}}/K.$$

2000 *Mathematics Subject Classification.* 32G20.

Key words and phrases. degenerating Hodge structure; period domain; toroidal compactification.

Now K is isomorphic to the unitary group. Then $G_{\mathbb{R}}/K$ can have a complex structure so that it is isomorphic to the Siegel space $\mathscr{H}$ or its complex conjugate $\bar{\mathscr{H}}$. This map is compatible with the action of $G_{\mathbb{Z}}$. Then we have the map

$$\Gamma\backslash D \to \Gamma\backslash\mathscr{H}, \quad \Gamma\backslash D \to \overline{\Gamma\backslash\mathscr{H}} \tag{1.1}$$

for a subgroup Γ of $G_{\mathbb{Z}}$.

In [**H**], we defined fans of even-type and odd-type. For an even-type (resp. odd-type) fan Σ which is compatible with Γ, we constructed the map

$$p^{\mathrm{ev}} : \Gamma\backslash D_{\Sigma} \to \Gamma\backslash\mathscr{H}_{\Sigma}, \quad (\text{resp. } p^{\mathrm{od}} : \Gamma\backslash D_{\Sigma} \to \overline{\Gamma\backslash\mathscr{H}}_{\Sigma})$$

as an extension of the map (1.1). Moreover we have the map $\zeta : \Gamma\backslash\mathscr{H}_{\Sigma} \to \Gamma\backslash\mathscr{H}_{S}$ to the Satake compactification $\Gamma\backslash\mathscr{H}_{S}$. We also showed that $\zeta\circ p^{\mathrm{ev}}$ and $\bar{\zeta}\circ p^{\mathrm{od}}$ are constructed by taking the boundary of the cycle space.

In this paper, we introduce a construction of fans of even-type and odd-type which is compatible with an arithmetic subgroup Γ. We show that the fans of even-type and odd-type are constructed by a decompositions of the open cones of the Lie group $\mathfrak{g}$ similar to the toroidal compactifications. Moreover, we define a subfan $\Sigma_{\mathrm{ev}} \subset \Sigma_{\mathrm{tor}}$ of even-type cones and prove the map

$$p^{\mathrm{ev}} : \Gamma\backslash D_{\Sigma_{\mathrm{ev}}} \to \Gamma\backslash\mathscr{H}_{\Sigma_{\mathrm{ev}}} (\subset \Gamma\backslash\mathscr{H}_{\Sigma_{\mathrm{tor}}})$$

is surjective.

In addition, recently Kerr and Pearlstein [**KP**] have given a construction of the partial compactifications of Mumford-Tate domains in the same manner as the Kato-Usui's construction. We expect that the construction of fans we discuss in this paper works in the case for Mumford-Tate domains.

2. Recipe for the fans

Let $H_{\mathbb{Z}}$ be a $\mathbb{Z}$-module of rank $2n$, and let Q be a bilinear form on $H_{\mathbb{Z}}$ defined by

$$\begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}.$$

Let $\{h^{p,-p-1}\}_p$ be a Hodge numbers of a Hodge structure of weight -1. Then we have a period domain D for $(H_{\mathbb{Z}}, Q, \{h^{p,-p-1}\}_p)$. Now

$$G_A = \mathrm{Aut}\,(H_A, Q) \cong Sp(n, A)$$

for $A = \mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$, and $G_{\mathbb{R}}$ acts on D transitively. The isotropy subgroup L and the maximal subgroup K including L are

$$L \cong \prod_p U(h^{p,-p-1}), \quad K \cong U(n).$$

We denote by $\mathscr{H}$ the Siegel space of degree n. Then $\mathscr{H} \cong G_{\mathbb{R}}/K$, and $\mathscr{H}$ is a period domain for

$$(H_{\mathbb{Z}}, Q, \{h^{0,-1} = h^{-1,0} = n,\ 0 \text{ otherwise}\}).$$

2.1. Polyhedral decompositions. Let S be an isotropic subspace of $H_{\mathbb{R}}$ with respect to Q. Now $0 \le \dim S \le n$ and S is corresponding to the Satake boundary component of $\mathscr{H}$ (cf. [**N**]). For the Lie algebra $\mathfrak{g}$ of $G_{\mathbb{R}}$, we define the subalgebra

$$\eta(S) = \{N \in \mathfrak{g} \mid \operatorname{Im} N \subset S\}.$$

For $X \in \mathfrak{g}$, we have $Q(Xv, w) + Q(v, Xw) = 0$ for $v, w \in H_{\mathbb{R}}$. Then $N^2 = 0$ for $N \in \eta(S)$ and all elements in $\eta(S)$ are commutative with each other. We define

$$\eta^+(S) = \{N \in \eta^+ \mid \phi_N > 0 \text{ on } H_{\mathbb{R}}/S^{\perp}\}$$

where ϕ_N is the bilinear form defined by $Q(\bullet, N\bullet)$. Let Γ be an arithmetic subgroup of $G_{\mathbb{Z}}$, and let $\Gamma(S)$ be the maximal subgroup of Γ which stabilizes S.

Definition 2.1. A fan $\Sigma(S)$ of cones[1] in the closure $\mathrm{Cl}\,(\eta^+(S))$ is said to be a $\Gamma(S)$-admissible polyhedral decomposition of $\eta^+(S)$ if:

- If $\sigma \in \Sigma(S)$ and $\gamma \in \Gamma(S)$, $\gamma\sigma \in \Sigma(S)$;
- There exists finitely many cones $\sigma_1, \dots, \sigma_\ell \in \Sigma(S)$ such that for any $\sigma \in \Sigma(S)$, there exists $\gamma \in \Gamma(S)$ with $\gamma\sigma = \sigma_j$ for some j;
- $\eta^+(S) = \bigcup_{\sigma \in \Sigma(S)} (\sigma \cap \eta^+(S))$.

Moreover a union $\Sigma = \bigcup_{Q(S,S)=0} \Sigma(S)$ of $\Gamma(S)$-admissible polyhedral decompositions is said to be a Γ-admissible family if:

- $\gamma\Sigma(S) = \Sigma(\gamma S)$;
- If $S' \subset S$, $\Sigma(S')$ is the restriction of $\Sigma(S)$ on $\mathrm{Cl}(\eta^+(S'))$.

The existence of a Γ-admissible family of polyhedral decompositions is proved by [**AMRT**].

2.2. Even-type and odd-type fans. Let σ be a nilpotent cone in $\mathfrak{g}$. σ can be written as

$$\sigma = \sum_{1 \le j \le \ell} \mathbb{R}_{\ge 0} N_j$$

with nilpotents $N_1, \dots, N_\ell \in \mathfrak{g}_{\mathbb{Q}}$. For F in the compact dual $\check{D}$, we call $(\sigma, \exp(\sigma_{\mathbb{C}})F)$ is a nilpotent orbit if it satisfies the following conditions:

- $\exp(\sum_j ix_j N_j)F \in D$ if $x_1, \dots, x_\ell \gg 0$;
- $NF^p \subset F^{p-1}$ for any p and $N \in \sigma$.

Let $(\sigma, \exp(\sigma_{\mathbb{C}})F)$ be a nilpotent orbit. We then have the weight filtration $W(\sigma)$ and $(W(\sigma), F)$ is a mixed Hodge structure twisting $W(\sigma)$. Now we have the Deligne decomposition $H_{\mathbb{C}} = \bigoplus_{p,q} I^{p,q}$ for $(W(\sigma), F)$ where

$$W(\sigma)_k = \sum_{p+q \le k} I^{p,q}, \quad F^\ell = \sum_{p \ge \ell} I^{p,q}.$$

Definition 2.2. A nilpotent orbit $(\sigma, \exp(\sigma_{\mathbb{C}})F)$ is called even-type (resp. odd-type) if it satisfies the following conditions:

- $N^2 = 0$ for N for $N \in \sigma$;
- $I^{p,-p} = 0$ for any odd (resp. even) integer p with respect to the LMHS $(W(\sigma), F)$.

---

[1] In this paper, all cones are assumed to be finitely generated convex rational polyhedral cone.

A nilpotent cone σ is called even-type (resp. odd-type) if σ generates a nilpotent orbit and all σ-nilpotent orbits are even-type (resp. odd-type). A fan Σ is called even-type (resp. odd-type) if any face of Σ is even-type (resp. odd-type).

Let $(\sigma, \exp(\sigma_{\mathbb{C}})F)$ be a nilpotent orbit. By the property of $W = W(\sigma)$ ([**S**, Lemma 6.4]),

$$W_{-2}^{\perp} = W_{-1}.$$

Then W_{-2} is an isotropy subspace. Moreover we have the following Lemma:

Lemma 2.3. *For the relative interior σ° of σ, $\sigma^{\circ} \subset \eta^{+}(W_{-2})$ (resp. $\sigma^{\circ} \subset -\eta^{+}(W_{-2})$) if, and only if σ is even-type (resp. odd-type).*

Proof. If σ is even-type (resp. odd-type), $N^2 = 0$ and then the weight filtration is given by

$$W(\sigma)_0 = H_{\mathbb{R}}, \quad W(\sigma)_{-1} = \operatorname{Ker} N, \quad W(\sigma)_{-2} = \operatorname{Im} N, \quad W(\sigma)_{-3} = \{0\}$$

where $N \in \sigma^{\circ}$. Therefore

$$H_{\mathbb{C}}/(W_{-2,\mathbb{C}})^{\perp} = H_{\mathbb{C}}/W_{-1,\mathbb{C}} \cong \bigoplus_{p:\text{even}} I^{p,-p} \quad (\text{resp. } \bigoplus_{p:\text{odd}} I^{p,-p}).$$

By the polarization condition of the LMHS,

$$i^{2p} Q(v, N\bar{v}) = \begin{cases} Q(v, N\bar{v}) > 0 & \text{if } p \text{ is even,} \\ -Q(v, N\bar{v}) > 0 & \text{if } p \text{ is odd,} \end{cases}$$

for $0 \neq v \in I^{p,-p}$ and $N \in \sigma^{\circ}$. Then $\sigma^{\circ} \subset \eta^{+}(W_{-2})$ (resp. $\sigma^{\circ} \subset -\eta^{+}(W_{-2})$), and vice versa. $\square$

2.3. Maps to the toroidal compactifications. Let $(\sigma, \exp(\sigma_{\mathbb{C}})F)$ be an even-type (resp. odd-type) nilpotent orbit. By [**CKS**, Proposition 2.20] there exists $\delta \in L_{\mathbb{R}}^{-1,-1}(W(\sigma), F)$ uniquely such that $(W(\sigma), e^{-i\delta}F)$ is $\mathbb{R}$-split LMHS. We write $\hat{F} = e^{-i\delta}F$. We then have the $\mathbb{R}$-split nilpotent orbit $(\sigma, \exp(\sigma_{\mathbb{C}})\hat{F})$ uniquely. Let $H_{\mathbb{C}} = \bigoplus_{p+q=0,-1,-2} I^{p,q}$ be the Deligne decomposition with respect to the LMHS $(W(\sigma), \hat{F})$. We define $\tilde{F} \in \check{\mathscr{H}}$ by

$$\tilde{F}^0 = \Big(\bigoplus_{p:\text{ even}} I^{p,-p-1}\Big) \oplus \Big(\bigoplus_{p} I^{p,-p}\Big),$$

$$(\text{resp. } \tilde{F}^0 = \Big(\bigoplus_{p:\text{ odd}} I^{p,-p-1}\Big) \oplus \Big(\bigoplus_{p} I^{p,-p}\Big))$$

Now $(W(\sigma), \tilde{F})$ (resp. $(W(\sigma), \overline{\tilde{F}})$) is a $\mathbb{R}$-split LMHS with the polarization $Q(\bullet, N^j \bullet)$ on Gr_W^j for $j \geq 0$ and $N \in \sigma^{\circ}$ (resp. $N \in -\sigma^{\circ}$). Then $(\sigma, \exp(\sigma_{\mathbb{C}})\tilde{F})$ is a $\mathbb{R}$-split nilpotent orbit for $\mathscr{H}$ (resp. $\bar{\mathscr{H}}$). By a linear algebraic calculation, we have the following proposition:

Proposition 2.4 ([**H**, Proposition 2.10]). $p^{\mathrm{ev}}(e^{zN}\hat{F}) = e^{zN}\tilde{F}$ *(resp.* $p^{\mathrm{od}}(e^{\bar{z}N}\hat{F}) = e^{\bar{z}N}\tilde{F}$*) for* $\operatorname{Im} z > 0$ *and* $N \in \sigma^{\circ}$.

We denote by $\mathbf{B}(\sigma)$ (resp. $\mathbf{B}_{\mathrm{tor}}(\sigma)$, $\bar{\mathbf{B}}_{\mathrm{tor}}(\sigma)$) the set of σ-nilpotent orbits for D (resp. $\mathscr{H}$, $\bar{\mathscr{H}}$). For an even-type (resp. odd-type) nilpotent cone σ, we define the map $p^{\mathrm{ev}} : \mathbf{B}(\sigma) \to \mathbf{B}_{\mathrm{tor}}(\sigma)$ (resp. $p^{\mathrm{od}} : \mathbf{B}(\sigma) \to \bar{\mathbf{B}}_{\mathrm{tor}}(\sigma)$) by

$$(\sigma, \exp(\sigma_{\mathbb{C}})F) \mapsto (\sigma, \exp(\sigma_{\mathbb{C}})e^{i\delta}\tilde{F})$$

where $F = e^{i\delta}\hat{F}$.

Proposition 2.5. *Let σ be an even-type cone such that $\sigma^\circ \subset \eta^+(S)$ with $\dim S \le h^{0,-1}$. Then $p^{\mathrm{ev}} : \mathbf{B}(\sigma) \to \mathbf{B}_{\mathrm{tor}}(\sigma)$ is surjective.*

Proof. For a nilpotent orbit $(\sigma, \exp(\sigma_{\mathbb{C}})F) \in \mathbf{B}_{\mathrm{tor}}(\sigma)$, it is enough to show that there exists a nilpotent orbit $(\sigma, \exp(\sigma_{\mathbb{C}})\breve{F}) \in \mathbf{B}(\sigma)$ such that the image through p^{ev} is $(\sigma, \exp(\sigma_{\mathbb{C}})F)$. Let $m = \dim S$. Then the Deligne decomposition for $(W(\sigma), F)$ is given by

$$H_{\mathbb{C}} = I^{0,0} \oplus I^{-1,0} \oplus I^{0,-1} \oplus I^{-1,-1}, \quad \dim I^{0,0} = m, \quad \dim I^{-1,0} = n-m.$$

Now $\overline{I^{-1,0}} = I^{0,-1}$, then $I^{-1,0} \oplus I^{0,-1}$ is defined over $\mathbb{R}$. Let H' be the subspace of $H_{\mathbb{R}}$ such that $H'_{\mathbb{C}} = I^{-1,0} \oplus I^{0,-1}$. Then $I^{-1,0}$ determines the point in the period domain for

$$(H', Q|_{H'}, \{h^{-1,0} = h^{0,-1} = n-m, \ 0 \text{ otherwise}\})$$

which is isomorphic to the Siegel space $\mathscr{H}'$ of degree $n-m$. Let D' be the period domain for $(H', Q|_{H'}, \{h'^{p,-1-p}\})$ where

$$h'^{p,-1-p} = \begin{cases} h^{p,-1-p} & \text{if } p \neq -1, 0, \\ h^{p,-1-p} - m & \text{if } p = -1, 0. \end{cases}$$

We then have

$$\mathscr{H}' \cong Sp(n-m, \mathbb{R})/U(n-m), \quad D' \cong Sp(n-m, \mathbb{R})/\prod_p U(h'^{p,-p-1}),$$

and we have the projection $\phi : D' \to \mathscr{H}'$. Now $I^{0,-1} \in \mathscr{H}'$ and there exists $F' \in D'$ such that $\phi(F') = I^{0,-1}$. We define $\breve{F} \in \check{D}$ by

$$\breve{F}^p = \begin{cases} F'^p & \text{if } p > 0 \\ F'^p \oplus I^{0,0} & \text{if } p = 0 \\ F'^p \oplus I^{0,0} \oplus I^{-1,-1} & \text{if } p < 0. \end{cases}$$

Then $(W(\sigma), \breve{F})$ is a LMHS such that

$$\mathrm{Gr}^{W(\sigma)}_{0,\mathbb{C}} \cong I^{0,0}, \quad \mathrm{Gr}^{W(\sigma)}_{-1}(\breve{F}) \cong F', \quad \mathrm{Gr}^{W(\sigma)}_{-2,\mathbb{C}} \cong I^{-1,-1}.$$

Therefore $(\sigma, \exp(\sigma_{\mathbb{C}})\breve{F})$ is a σ-nilpotent orbit for D and the image through p^{ev} is $(\sigma, \exp(\sigma_{\mathbb{C}})F)$. □

Remark 2.6. The above proposition does not hold for an odd-type cone. See the example of §2.4.

For a fan Σ of nilpotent cones, we define

$$D_\Sigma = \bigsqcup_{\sigma \in \Sigma} \mathbf{B}(\sigma).$$

Then for an even-type (resp. odd-type) fan Σ we can define the map

$$p^{\mathrm{ev}} : D_\Sigma \to \mathscr{H}_\Sigma \quad (\text{resp. } p^{\mathrm{od}} : D_\Sigma \to \bar{\mathscr{H}}_\Sigma),$$

and for a subgroup Γ of $G_{\mathbb{Z}}$ which is compatible with Σ we have

$$p^{\mathrm{ev}} : \Gamma\backslash D_\Sigma \to \Gamma\backslash \mathscr{H}_\Sigma \quad (\text{resp. } p^{\mathrm{od}} : \Gamma\backslash D_\Sigma \to \overline{\Gamma\backslash \mathscr{H}}_\Sigma),$$

where $p^{\mathrm{ev}} = id$ if $D = \mathscr{H}$.

Let Γ be an arithmetic subgroup of $G_{\mathbb{R}}$ and let Σ_{tor} be a Γ-admissible family of polyhedral decompositions. Then $\Gamma\backslash D_{\Sigma_{\mathrm{tor}}}$ is called a toroidal compactification.

A toroidal compactification is compact and moreover is smooth if Γ is neat. By Lemma 2.3, for $\sigma \in \Sigma_{\mathrm{tor}}$, σ is even-type and $-\sigma$ is odd-type if it has nilpotent orbit. We then have the map

$$p^{\mathrm{ev}} : \Gamma\backslash D_{\Sigma_{\mathrm{tor}}} \to \Gamma\backslash\mathscr{H}_{\Sigma_{\mathrm{tor}}}, \quad p^{\mathrm{od}} : \Gamma\backslash D_{\Sigma^-_{\mathrm{tor}}} \to \overline{\Gamma\backslash\mathscr{H}}_{\Sigma^-_{\mathrm{tor}}} \tag{2.1}$$

where Σ^-_{tor} is the fan of $-\sigma$ for all $\sigma \in \Sigma_{\mathrm{tor}}$. As we will see at §2.4, these maps may not be surjective.

Now

$$\Sigma_{\mathrm{tor}} = \bigcup_{Q(S,S)=0} \Sigma(S).$$

We then define the subfan

$$\Sigma_{\mathrm{ev}} = \bigcup_{\substack{Q(S,S)=0,\\ \dim S \le h^{0,-1}}} \Sigma(S) \tag{2.2}$$

of Σ_{tor}. By Proposition 2.5, we have the following corollary

Corollary 2.7. *$p^{\mathrm{ev}} : \Gamma\backslash D_{\Sigma_{\mathrm{ev}}} \to \Gamma\backslash\mathscr{H}_{\Sigma_{\mathrm{ev}}}$ is surjective.*

2.4. Example. We consider the case where $h^{p,-p-1} = 1$ if $p = 1, 0, -1, -2$, $h^{p,-p-1} = 0$ otherwise (the case for Hodge structures of Calabi-Yau threefolds with $h^{2,1} = 1$). In this case

$$G_{\mathbb{R}} \cong Sp(2,\mathbb{R}), \quad L \cong U(1)\times U(1), \quad K \cong U(2).$$

In [**KU**, §12.3], this case is well-studied. Any nilpotent cone in this case is rank 1, and its generator N is classified as follows:

(I) $N^2 = 0,\ \dim(\operatorname{Im} N) = 1$;
(II) $N^2 = 0,\ \dim(\operatorname{Im} N) = 2$;
(III) $N^3 \neq 0, N^4 = 0$.

LMHS of type-I and type-II are described as follows:



(I)



(II)

Then type-I cone is even-type and type-II cone is odd-type. The fan of all possible type-I cones is

$$\Sigma_1 = \{\mathbb{R}_{\ge 0}N \mid N \in \mathfrak{g}_{\mathbb{Q}},\ N^2 = 0,\ \dim(\operatorname{Im} N) = 1,\ \phi_N > 0 \text{ on } H_{\mathbb{R}}/\operatorname{Ker} N\}.$$

For a generator N of $\sigma \in \Sigma_1$, $\eta(\operatorname{Im} N)$ is 1-dimensional, and then $\eta^+(\operatorname{Im} N) = \mathbb{R}_{>0}N$. Then $\Sigma_1 = \Sigma_{\mathrm{ev}}$ of (2.2).

On the other hand, the fan Σ_2 of all possible type-II cones, all cones of Σ_2 are rank 1. For a generator N of $\sigma \in \Sigma_2$, $\dim(\operatorname{Im} N) = 2$ and $\dim(\eta(\operatorname{Im} N)) = 3$. Then Σ_2 does not contain a polyhedral decomposition of $\eta^+(\operatorname{Im} N)$.

For a type-I cone σ_1, by Proposition 2.5 the map

$$p^{\mathrm{ev}} : \mathbf{B}(\sigma_1) \to \mathbf{B}_{\mathrm{tor}}(\sigma_1)$$

is surjective. In fact,

$$\mathbf{B}(\sigma_1) \cong \bar{\mathscr{H}} \times \mathbb{C}, \quad \mathbf{B}_{\mathrm{tor}}(\sigma_1) \cong \mathscr{H} \times \mathbb{C}.$$

The map p^{ev} on the boundary component is given by the complex conjugation of the first element.

For a type-II cone σ_2, we have $\mathbf{B}(\sigma_2) \cong \mathbb{C} \times \{\pm 1\}$ by [**KU**, §12.3]. Then

$$\dim \mathbf{B}(\sigma_2) < \dim \bar{\mathbf{B}}_{\mathrm{tor}}(\sigma_2) = 2.$$

Therefore the map p^{od} on the boundary component is not surjective.

References

[AMRT] A. Ash, D. Mumford, M. Rapoport and Y. S. Tai, *Smooth compactification of locally symmetric varieties,* Math. Sci. Press, Brookline, 1975.

[CCK] J. A. Carlson, E. Cattani and A. Kaplan, *Mixed Hodge structures and compactification of Siegel's space*, in Journées de géométrie algébrique d'Angers 1979 (A. Beauville, ed.), Sijthohoff & Nordhoff, 1980, 77–105.

[CKS] E. Cattani, A. Kaplan and W. Schmid, *Degeneration of Hodge structures*, Ann. of Math. **123** (1986), 457–535.

[G] P. Griffiths, *Periods of integrals on algebraic manifolds. I. Construction and properties of the modular varieties*, Amer. J. Math. **90** (1968) 568–626.

[H] T. Hayama, *Boundaries of cycle spaces and degenerating Hodge structures*, arXiv:1203.6770

[KU] K. Kato and S. Usui, *Classifying space of degenerating polarized Hodge structures*, Annals of Mathematics Studies, **169**. Princeton University Press, Princeton, NJ, 2009.

[KP] M. Kerr and G. Pearlstein, *Boundary components of Mumford-Tate domains*, preprint.

[N] Y. Namikawa, *Toroidal compactification of Siegel spaces*, Lecture Notes in Math **812**, Springer-Verlag, 1980.

[S] W. Schmid, *Variation of Hodge structure: the singularities of the period mapping*, Invent. Math. **22** (1973), 211–319.

Mathematical Sciences Center, Tsinghua University, Haidian District, Beijing 100084, China

E-mail address: `tatsuki@math.tsinghua.edu.cn`